ACTIVE MICRO TECHNOLOGIES

107 Technology Drive • Lincolnton, NC 28092
(704) 276-7100 • Fax (704) 276-7101

OECD TG 442C: *In Chemico* Skin Sensitization

Tradename: Leucidal® SF Max Powder

Code: M15031

CAS #: 1686112-36-6 (or) 68333-16-4

Test Request Form #: 10288

Lot #: 841500

Sponsor: *Active Micro Technologies, LLC; 107 Technology Drive Lincolnton, NC 28092*
Study Director: *Maureen Drumwright*
Principle Investigator: *Daniel Shill*

Test Performed:
OECD TG 442C: *In Chemico* Skin Sensitization
Direct Peptide Reactivity Assay (DPRA)

Introduction

A skin sensitizer is a substance that will lead to an allergic response following skin contact[1]. Haptenation is the covalent binding of a hapten, or low-molecular weight substance or chemical, to proteins in the skin. This is considered the prominent mechanism which defines a chemical as a sensitizer. Haptenation is described as a "molecular initiating event" in the OECD Adverse Outcome Pathway (AOP) for skin sensitization which summarizes the key events known to be involved in chemically-induced allergic contact dermatitis[2]. The direct peptide reactivity assay (DPRA) is designed to mimic the covalent binding of electrophilic chemicals to nucleophilic centers in skin proteins by quantifying the reactivity of chemicals towards the model synthetic peptides containing cysteine and lysine. The DPRA is able to distinguish sensitizers from non-sensitizer with 82% accuracy (sensitivity of 76%; specificity of 92%)[3].

This assay was conducted to determine skin sensitization hazard of **Leucidal® SF Max Powder** in accordance with European Union Reference Laboratory for Alternatives to Animal Testing (EURL ECVAM) and OECD Test Guideline 442C.

Assay Principle

The DPRA is an *in chemico* method which addresses peptide reactivity by measuring depletion of synthetic heptapeptides containing either cysteine or lysine following 24 hours incubation with the test substance. The peptide is a custom material containing phenylalanine to aid in detection. Depletion of the peptide in the reaction mixture is measured by HPLC with gradient elution and UV detection at 220 nm. Cysteine and lysine peptide percent depletion values are then calculated and used in a prediction model which allows assigning the test chemical to one of four reactivity classes used to support the discrimination between sensitizers and non-sensitizers.

1. United Nations Economic Commission (UNECE) (2013) Global Harmonized System of Classification and Labelling of Chemicals (GHS) 5th Revised Edition
2. OECD (2012). The Adverse Outcome Pathway for Skin Sensitization Initiated by Covalent Binding to Proteins. Part 1: Scientific Evidence. Series on Testing and Assessment No. 168
3. EC EURL ECVAM (2012) Direct peptide reactivity assay (DPRA) validation study report; pp 1 -74.

Information contained in this technical literature is believed to be accurate and is offered in good faith for the benefit of the customer. The company, however, cannot assume any liability or risk involved in the use of its chemical products since the conditions of use are beyond our control. Statements concerning the possible use of our products are not intended as recommendations to use our products in the infringement of any patent. We make no warranty of any kind, expressed or implied, other than that the material conforms to the applicable standard specification.

107 Technology Drive • Lincolnton, NC 28092
(704) 276-7100 • Fax (704) 276-7101

Materials

A.	**Equipment:**	HPLC-UV (Waters Breeze - Waters 2998 Photodiode Array Detector); Pipettes; Analytical balance
B.	**HPLC/Guard Columns:**	Agilent Zorbax SB-C18 2.1mm x 100mm x 3.5µm; Phenomenex Security Guard C18 4mm x 2mm
C.	**Chemicals:**	Trifluoroacetic acid; Ammonium acetate; Ammonium hydroxide; Acetonitrile; Cysteine peptide (Ac-RFAA**C**AA-COOH); Lysine peptide (Ac-RFAA**K**AA-COOH); Cinnamic aldehyde
D.	**Reagents/Buffers:**	Sodium phosphate buffer (100mM); Ammonium acetate buffer (100mM)
E.	**Other:**	Sterile disposable pipette tips

Methods

Solution Preparation:

- 0.667mM Cysteine Peptide in 100mM Phosphate Buffer (pH 7.5)
- 0.667mM Lysine Peptide in 100mM Ammonium Acetate Buffer (pH 10.2)
- 100mM Cinnamic Aldehyde in Acetonitrile
- 100mM* **Leucidal® SF Max Powder** in Acetonitrile

*For mixtures and multi-constituent substances of known composition such as **Leucidal® SF Max Powder**, a single purity should be determined by the sum of the proportion of its constituents (excluding water), and a single apparent molecular weight determined by considering the individual molecular weights of each component in the mixture (excluding water) and their individual proportions. The resulting purity and apparent molecular weight can then be used to calculate the weight of test chemical necessary to prepare a 100 mM solution.

Reference Controls:

- Reference Control A: For calibration curve accuracy
- Reference Control B: For peptide stability over analysis time of experiment
- Reference Control C: For verification that the solvent does not impact percent peptide depletion

Sample, Reference Control, and Co-Elution Control Preparation:

- Once these solutions have been made they should be incubated at room temperature, protected from light, for 24±2 hours before running HPLC analysis.
- Each chemical should be analyzed in triplicate.

1:10 Ratio, Cysteine Peptide 0.5mM Peptide, 5mM Test Chemical	1:50 Ratio, Lysine Peptide 0.5mM Peptide, 25mM Test Chemical
• 750µL Cysteine Peptide Solution (or 100mM Phosphate Buffer, pH 7.5, for Co-Elution Controls) • 200µL Acetonitrile • 50µL Test Chemical Solution (or Acetonitrile for Reference Controls)	• 750µL Lysine Peptide Solution (or 100mM Ammonium Acetate Buffer, pH 10.2, for Co-Elution Controls) • 250µL Test Chemical Solution (or Acetonitrile for Reference Controls)

Information contained in this technical literature is believed to be accurate and is offered in good faith for the benefit of the customer. The company, however, cannot assume any liability or risk involved in the use of its chemical products since the conditions of use are beyond our control. Statements concerning the possible use of our products are not intended as recommendations to use our products in the infringement of any patent. We make no warranty of any kind, expressed or implied, other than that the material conforms to the applicable standard specification.

107 Technology Drive • Lincolnton, NC 28092
(704) 276-7100 • Fax (704) 276-7101

Calibration Curve:

- Standards are prepared in a solution of 20% Acetonitrile:Buffer
  - For the Cysteine peptide using the phosphate buffer, pH 7.5
  - For the Lysine peptide using the ammonium acetate buffer, pH 10.2

	Standard 1	Standard 2	Standard 3	Standard 4	Standard 5	Standard 6	Standard 7
mM Peptide	0.534	0.267	0.1335	0.0667	0.0334	0.0167	0.000

HPLC Analysis:

- HPLC-UV system should be equilibrated at 30°C with 50% Mobile Phase A (0.1% (v/v) trifluoroacetic acid in water) and 50% Mobile Phase B (0.085% (v/v) trifluoroacetic acid in acetonitrile) for 2 hours
- Absorbance is measured at 220nm
- Flow Conditions:

Time	Flow	%A	%B
0 minutes	0.35 mL/min	90	10
10 minutes	0.35 mL/min	75	25
11 minutes	0.35 mL/min	10	90
13 minutes	0.35 mL/min	10	90
13.5 minutes	0.35 mL/min	90	10
20 minutes	End Run		

Data and Reporting

Acceptance Criteria:

1. The following criteria must be met for a run to be considered valid:
   a. Standard calibration curve should have an $r^2 > 0.99$.
   b. Mean percent peptide depletion values of three replicates for the positive control cinnamic aldehyde should be between 60.8% and 100% for the cysteine peptide and between 40.2% and 69% for the lysine peptide and the maximum standard deviation should be <14.9 for the percent cysteine depletion and <11.6 for the percent lysine depletion.
   c. Mean peptide concentration of reference controls A should be 0.50±0.05mM and the coefficient of variable of the peptide peak areas for reference B and C in acetonitrile should be <15.0%.

2. The following criteria must be met for a test chemical's results to be considered valid:
   a. Maximum standard deviation should be <14.9 for percent cysteine depletion and <11.6 for percent lysine depletion.
   b. Mean peptide concentration of the three reference control C should be 0.50±0.05mM.

Information contained in this technical literature is believed to be accurate and is offered in good faith for the benefit of the customer. The company, however, cannot assume any liability or risk involved in the use of its chemical products since the conditions of use are beyond our control. Statements concerning the possible use of our products are not intended as recommendations to use our products in the infringement of any patent. We make no warranty of any kind, expressed or implied, other than that the material conforms to the applicable standard specification.

107 Technology Drive • Lincolnton, NC 28092
(704) 276-7100 • Fax (704) 276-7101

Prediction Model:

Cysteine 1:10/Lysine 1:50 Prediction Model		
Mean of Cysteine and Lysine % Depletion	**Reactivity Class**	**Prediction**
0% < Mean % Depletion < 6.38%	Minimal Reactivity	Non-sensitizer
6.38% < Mean % Depletion < 22.62%	Low Reactivity	Sensitizer
22.62% < Mean % Depletion < 42.47%	Moderate Reactivity	Sensitizer
42.47% < Mean % Depletion < 100%	High Reactivity	Sensitizer

If co-elution occurs with the lysine peptide, than the cysteine 1:10 prediction model can be used:

Cysteine 1:10 Prediction Model		
Mean of Cysteine and Lysine % Depletion	**Reactivity Class**	**Prediction**
0% < Cys % Depletion < 13.89%	Minimal Reactivity	Non-sensitizer
13.89% < Cys % Depletion < 23.09%	Low Reactivity	Sensitizer
23.09% < Cys % Depletion < 98.24%	Moderate Reactivity	Sensitizer
98.24% < Cys % Depletion < 100%	High Reactivity	Sensitizer

Therefore the measured values of % depletion in the three separated runs for each peptide depletion assay include:

Cysteine 1:10/Lysine 1:50 Prediction Model		
Mean of Cysteine and Lysine % Depletion	**Reactivity Class**	**Prediction**
3.09	Minimal Reactivity	Non-sensitizer
3.13	Minimal Reactivity	Non-sensitizer
3.14	Minimal Reactivity	Non-sensitizer

Cysteine 1:10 Prediction Model		
Mean of Cysteine and Lysine % Depletion	**Reactivity Class**	**Prediction**
2.99	Minimal Reactivity	Non-sensitizer
3.03	Minimal Reactivity	Non-sensitizer
3.02	Minimal Reactivity	Non-sensitizer

Results and Discussion

The data obtained from this study met criteria for a valid assay and the controls performed as anticipated.

Percent peptide depletion is determined by the following equation:

$$Percent\ Peptide\ Depletion = \left[1 - \left(\frac{Peptide\ Peak\ Area\ in\ Replicate\ Injection}{Mean\ Peptide\ Peak\ Area\ in\ Reference\ Controls\ C}\right)\right] \times 100$$

Based on HPLC-UV analysis of **Leucidal® SF Max Powder (M15031)** we can determine this product is not classified as a sensitizer and is not predicted to cause allergic contact dermatitis. The Mean Percent Depletion of Cysteine and Lysine was 3.07% causing minimal reactivity in the assay giving us the prediction of a non-sensitizer.

Information contained in this technical literature is believed to be accurate and is offered in good faith for the benefit of the customer. The company, however, cannot assume any liability or risk involved in the use of its chemical products since the conditions of use are beyond our control. Statements concerning the possible use of our products are not intended as recommendations to use our products in the infringement of any patent. We make no warranty of any kind, expressed or implied, other than that the material conforms to the applicable standard specification.